

ISAS/94/91/EP

NORDITA 91/47

On the S-matrix of the Sub-leading Magnetic Deformation of the Tricritical Ising Model in Two Dimensions

F. Colomo^{1*}, A. Koubek², G. Mussardo^{1†}

¹*NORDITA, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

²*International School for Advanced Studies, Strada Costiera 11, 34014 Trieste, Italy*

Abstract

We compute the S -matrix of the Tricritical Ising Model perturbed by the sub-leading magnetic operator using Smirnov's RSOS reduction of the Izergin-Korepin model. The massive model contains kink excitations which interpolate between two degenerate asymmetric vacua. As a consequence of the different structure of the two vacua, the crossing symmetry is implemented in a non-trivial way. We use finite-size techniques to compare our results with the numerical data obtained by the Truncated Conformal Space Approach and find good agreement.

June 1991

*Angelo Della Riccia Foundation's fellow.

†On leave of absence from: International School for Advanced Studies, Strada Costiera 11, 34014 Trieste, Italy.

1 Introduction

It has been pointed out by Zamolodchikov that certain deformations of minimal models of Conformal Field Theories (CFT) produce integrable massive field theories, which are characterized on mass-shell by a factorizable S -matrix [1]. A particularly interesting situation occurs for the “Tricritical Ising Model” (TIM) at the fixed point perturbed by the operator $\Phi_{2,1}$. This field has anomalous dimensions $(\Delta, \overline{\Delta}) = (\frac{7}{16}, \frac{7}{16})$ and is identified with the sub-leading magnetic operator¹, odd with respect to the Z_2 spin-reversal transformation. Hence, this deformation explicitly breaks the Z_2 symmetry of the tricritical point and the corresponding massive theory can exhibit “ Φ^3 -property”, i.e. the absence of a conserved current of spin 3 and the possibility to form a bound state through the process $AA \rightarrow A \rightarrow AA$. The counting argument supports this picture, giving for the spin of the conserved currents the values $s=(1,5,7,11,13)$ [1, 2]. The interesting features of this massive field theory have been first outlined in ref. [2] where the model was studied by the “Truncated Conformal Space Approach” (TCSA), proposed in ref. [3]. This approach consists in the diagonalization of the perturbed Hamiltonian

$$H = H_{CFT} + \lambda \int \Phi_{2,1}(x) dx \quad (1.1)$$

on a strip² of width R , truncated at a certain level of the Hilbert space defined by the conformal field theory at the fixed point. The lowest energy levels are given in fig. 1. The theory presents two degenerate ground states (which correspond to the minima of the asymmetric double-well Landau-Ginzburg potential in fig. 2) and a single excitation B of mass m below the threshold at $2m$. The double degeneracy of the vacuum permits two fundamental kink configurations $|K_+\rangle$ and $|K_-\rangle$ and, possibly, bound states thereof. If the two vacua were related by a symmetry transformation, one would expect a double degeneracy of the breather-like bound state $|B\rangle$ in the infrared regime $R \rightarrow \infty$. However, the absence of a Z_2 symmetry makes it possible that in this case only one of the two asymptotic states $|K_+K_-\rangle$ or $|K_-K_+\rangle$ is coupled to the bound state $|B\rangle$.

¹In the Landau-Ginzburg theory, the TIM represents the universality class of the φ^6 model. The operator $\Phi_{2,1}$ is represented by the third-order field φ^3 .

²We consider in the following only the case of periodic boundary condition.

A scattering theory for such model with asymptotic states $|K_+\rangle$, $|K_-\rangle$ and $|B\rangle$ was first conjectured in [4]. On the other hand, a general framework for the $\Phi_{1,2}$ and $\Phi_{2,1}$ deformations of CFT has been recently proposed in [5]. It is based on the RSOS reduction of the Izergin-Korepin model [6]. In section 3 we explicitly work out the RSOS-like S -matrix in the case of the TIM perturbed by the subleading magnetic operator. The scattering theory we obtain is different from the one of ref. [4], although both of them have the following features:

1. the existence of two fundamental kink configurations with mass m ;
2. the appearance of only one bound state of the above, with the same mass m .

Hence, both scattering theories give rise to a picture that qualitatively agrees with the results of ref. [2]. Therefore a more detailed analysis is required in order to decide which of the two theories is appropriate for the description of the scaling region of the TIM in presence of a third-order magnetic perturbation. As suggested in ref. [4], possible insight comes from the study of finite-size effects for the theory defined on the cylinder [8]. These finite-size effects, which can be directly related to the scattering data, control the exponential decay of the one-particle energy level to its asymptotic value.

The paper is organized as follows: in sect. 2 we outline Smirnov's RSOS reduction of the Izergin-Korepin model. In sect. 3 we present our results on the S -matrix of the $\Phi_{2,1}$ perturbation of TIM. Sect. 4 contains the analysis of the phase shifts and the asymptotic behaviour of the RSOS S -matrix. Sect. 5 consists of a short summary of Zamolodchikov's proposal for the S -matrix [4]. This proposal, together with ours, will be checked against numerical data in sect. 6, where we perform an accurate analysis of the energy levels obtained from the TCSA using the finite-size theory. Our conclusions are in sect. 7.

2 Smirnov's RSOS reduction of the Izergin-Korepin model

In a recent paper, Smirnov has related the $\Phi_{1,2}$ and $\Phi_{2,1}$ deformations of a CFT to a RSOS projection of the R -matrix of the Izergin-Korepin (IK) model [5, 6]. His results can be

summarized as follows. The R -matrix of the IK model cannot be directly interpreted as the S -matrix of the massive excitations of the perturbed CFT because, introducing a Hilbert space structure, it does not satisfy the unitarity requirement. But it is possible to use the quantum $SL(2)_q$ symmetry of the IK model to study the RSOS restriction of the Hilbert space. This happens when q is a p th root of unity. The RSOS reduction preserves the locality of an invariant set of operators and yields S -matrices which have a sensible physical interpretation. The RSOS states appearing in the reduced model

$$| \beta_1, j_1, k_1, | a_1 | \beta_2, j_2, k_2, \dots | a_{n-1} | \beta_n, j_n, k_n \rangle \quad (2.1)$$

are characterized by their rapidity β_i , by their type k (which distinguish the kinks from the breathers), by their $SL(2)_q$ spin j and by a string of integer numbers a_i , satisfying

$$a_i \leq \frac{p-2}{2} \ , \quad | a_k - 1 | \leq a_{k+1} \leq \min(a_k + 1, p - 3 - a_k) \ . \quad (2.2)$$

In the case of $\Phi_{2,1}$ deformation, using the following parameterization of the central charge of the original CFT

$$c = 1 - 6 \left(\frac{\pi}{\gamma} + \frac{\gamma}{\pi} - 2 \right) \ , \quad (2.3)$$

the q -parameter is given by

$$q = \exp(2i\gamma) \ . \quad (2.4)$$

For the unitary minimal CFT, $\gamma = \pi \frac{r}{r+1}$ ($r = 3, 4, \dots$). It is also convenient to define the quantity

$$\xi = \frac{2}{3} \left(\frac{\pi^2}{2\gamma - \pi} \right) \ . \quad (2.5)$$

The S -matrix of the RSOS states is given by [5]

$$\begin{aligned}
S_{a_{k-1}a_{k+1}}^{a_k a'_k}(\beta_k - \beta_{k+1}) = & \\
& \frac{i}{4} S_0(\beta_k - \beta_{k+1}) \left[\left\{ \begin{array}{ccc} 1 & a_{k-1} & a_k \\ 1 & a_{k+1} & a'_k \end{array} \right\}_q \right. \\
& \times \left(\left(\exp \left(\frac{2\pi}{\xi} (\beta_{k+1} - \beta_k) \right) - 1 \right) q^{(c_{a_{k+1}} + c_{a_{k-1}} - c_{a_k} - c_{a'_k} + 3)/2} \right. \\
& \left. - \left(\exp \left(-\frac{2\pi}{\xi} (\beta_{k+1} - \beta_k) \right) - 1 \right) q^{-(c_{a_{k+1}} + c_{a_{k-1}} - c_{a_k} - c_{a'_k} + 3)/2} \right) \\
& \left. + q^{-5/2} (q^3 + 1) (q^2 - 1) \delta_{a_k, a'_k} \right]. \tag{2.6}
\end{aligned}$$

Herein, c_a are the Casimir of the representation a , $c_a = a(a+1)$ and the expression of the $6j$ -symbols is that in ref. [7]. $S_0(\beta)$ has the following integral representation

$$\begin{aligned}
S_0(\beta) = & \left(\sinh \frac{\pi}{\xi} (\beta - i\pi) \sinh \frac{\pi}{\xi} \left(\beta - \frac{2\pi i}{3} \right) \right)^{-1} \\
& \times \exp \left(-2i \int_0^\infty \frac{dx}{x} \frac{\sin \beta x \sinh \frac{\pi x}{3} \cosh \left(\frac{\pi}{6} - \frac{\xi}{2} \right) x}{\cosh \frac{\pi x}{2} \sinh \frac{\xi x}{2}} \right). \tag{2.7}
\end{aligned}$$

3 RSOS S -matrix of $\Phi_{2,1}$ perturbed TIM

In the TIM perturbed by the subleading magnetization operator, $r = 4$ and $\xi = \frac{10\pi}{9}$. From eq. (2.2), the only possible values of a_i are 0 and 1 and the one-particle states are the vectors: $|K_{01}\rangle$, $|K_{10}\rangle$ and $|K_{11}\rangle$. All of them have the same mass m . Notice that the state $|K_{00}\rangle$ is not allowed. A basis for the two-particle asymptotic states is

$$|K_{01}K_{10}\rangle, |K_{01}K_{11}\rangle, |K_{11}K_{11}\rangle, |K_{11}K_{10}\rangle, |K_{10}K_{01}\rangle. \tag{3.1}$$

The scattering processes are

$$\begin{aligned}
|K_{01}(\beta_1)K_{10}(\beta_2)\rangle &= S_{00}^{11}(\beta_1 - \beta_2) |K_{01}(\beta_2)K_{10}(\beta_1)\rangle \\
|K_{01}(\beta_1)K_{11}(\beta_2)\rangle &= S_{01}^{11}(\beta_1 - \beta_2) |K_{01}(\beta_2)K_{11}(\beta_1)\rangle \\
|K_{11}(\beta_1)K_{10}(\beta_2)\rangle &= S_{10}^{11}(\beta_1 - \beta_2) |K_{11}(\beta_2)K_{10}(\beta_1)\rangle \\
|K_{11}(\beta_1)K_{11}(\beta_2)\rangle &= S_{11}^{11}(\beta_1 - \beta_2) |K_{11}(\beta_2)K_{11}(\beta_1)\rangle + S_{11}^{10}(\beta_1 - \beta_2) |K_{10}(\beta_2)K_{01}(\beta_1)\rangle \\
|K_{10}(\beta_1)K_{01}(\beta_2)\rangle &= S_{11}^{00}(\beta_1 - \beta_2) |K_{10}(\beta_2)K_{01}(\beta_1)\rangle + S_{11}^{10}(\beta_1 - \beta_2) |K_{11}(\beta_2)K_{11}(\beta_1)\rangle
\end{aligned} \tag{3.2}$$

Explicitly, the above amplitudes are given by

$$\begin{array}{c} 1 \\ \diagdown \quad \diagup \\ 0 \quad 0 \\ \diagup \quad \diagdown \\ 1 \end{array} = S_{00}^{11}(\beta) = \frac{i}{2} S_0(\beta) \sinh\left(\frac{9}{5}\beta - i\frac{\pi}{5}\right) \quad (3.3.a)$$

$$\begin{array}{c} 1 \\ \diagdown \quad \diagup \\ 0 \quad 1 \\ \diagup \quad \diagdown \\ 1 \end{array} = S_{01}^{11}(\beta) = -\frac{i}{2} S_0(\beta) \sinh\left(\frac{9}{5}\beta + i\frac{\pi}{5}\right) \quad (3.3.b)$$

$$\begin{array}{c} 1 \\ \diagdown \quad \diagup \\ 1 \quad 1 \\ \diagup \quad \diagdown \\ 1 \end{array} = S_{11}^{11}(\beta) = \frac{i}{2} S_0(\beta) \frac{\sin\left(\frac{\pi}{5}\right)}{\sin\left(\frac{2\pi}{5}\right)} \sinh\left(\frac{9}{5}\beta - i\frac{2\pi}{5}\right) \quad (3.3.c)$$

$$\begin{array}{c} 1 \\ \diagdown \quad \diagup \\ 1 \quad 1 \\ \diagup \quad \diagdown \\ 0 \end{array} = S_{11}^{01}(\beta) = -\frac{i}{2} S_0(\beta) \left(\frac{\sin\left(\frac{\pi}{5}\right)}{\sin\left(\frac{2\pi}{5}\right)}\right)^{\frac{1}{2}} \sinh\left(\frac{9}{5}\beta\right) \quad (3.3.d)$$

$$\begin{array}{c} 0 \\ \diagdown \quad \diagup \\ 1 \quad 1 \\ \diagup \quad \diagdown \\ 0 \end{array} = S_{11}^{00}(\beta) = -\frac{i}{2} S_0(\beta) \frac{\sin\left(\frac{\pi}{5}\right)}{\sin\left(\frac{2\pi}{5}\right)} \sinh\left(\frac{9}{5}\beta + i\frac{2\pi}{5}\right) \quad (3.3.e)$$

The function $S_0(\beta)$ can actually be computed. It is given by

$$\begin{aligned} S_0(\beta) &= -\left(\sinh \frac{9}{10}(\beta - i\pi) \sinh \frac{9}{10}\left(\beta - \frac{2\pi i}{3}\right)\right)^{-1} \\ &\quad \times w\left(\beta, -\frac{1}{5}\right) w\left(\beta, +\frac{1}{10}\right) w\left(\beta, \frac{3}{10}\right) \\ &\quad \times t\left(\beta, \frac{2}{9}\right) t\left(\beta, -\frac{8}{9}\right) t\left(\beta, \frac{7}{9}\right) t\left(\beta, -\frac{1}{9}\right), \end{aligned} \quad (3.4)$$

where

$$\begin{aligned} w(\beta, x) &= \frac{\sinh\left(\frac{9}{10}\beta + i\pi x\right)}{\sinh\left(\frac{9}{10}\beta - i\pi x\right)}; \\ t(\beta, x) &= \frac{\sinh\frac{1}{2}(\beta + i\pi x)}{\sinh\frac{1}{2}(\beta - i\pi x)}. \end{aligned}$$

It is easy to check the unitarity equations:

$$S_{11}^{00}(\beta) S_{11}^{00}(-\beta) + S_{11}^{01}(\beta) S_{11}^{10}(-\beta) = 1;$$

$$\begin{aligned}
S_{11}^{10}(\beta) S_{11}^{01}(-\beta) + S_{11}^{11}(\beta) S_{11}^{11}(-\beta) &= 1 ; \\
S_{11}^{10}(\beta) S_{11}^{00}(-\beta) + S_{11}^{11}(\beta) S_{11}^{10}(-\beta) &= 0 ; \\
S_{10}^{11}(\beta) S_{10}^{11}(-\beta) &= 1 ; \\
S_{00}^{11}(\beta) S_{00}^{11}(-\beta) &= 1 .
\end{aligned} \tag{3.5}$$

An interesting property of this S matrices is that the crossing symmetry occurs in a non-trivial way, i.e.

$$\begin{aligned}
S_{11}^{11}(i\pi - \beta) &= S_{11}^{11}(\beta) ; \\
S_{11}^{00}(i\pi - \beta) &= a^2 S_{00}^{11}(\beta) ; \\
S_{11}^{01}(i\pi - \beta) &= a S_{01}^{11}(\beta) ;
\end{aligned} \tag{3.6}$$

where

$$a = - \left(\frac{s\left(\frac{1}{5}\right)}{s\left(\frac{2}{5}\right)} \right)^{\frac{1}{2}}, \tag{3.7}$$

and $s(x) \equiv \sin(\pi x)$.

The above crossing-symmetry relations may be seen as due to a non-trivial charge conjugation operator (see also [9]). In most cases, the charge conjugation is implemented trivially, i.e. with $a = \pm 1$ in eq. (3.6). Here, the asymmetric Landau-Ginzburg potential distinguishes between the two vacua and gives rise to the value (3.7).

The amplitudes (3.3) are periodic along the imaginary axis of β with period $10\pi i$. The whole structure of poles and zeros is quite rich. On the physical sheet, $0 \leq \text{Im } \beta \leq i\pi$, the poles of the S -matrix are located at $\beta = \frac{2\pi i}{3}$ and $\beta = \frac{i\pi}{3}$ (fig. 3). The first pole corresponds to a bound state in the direct channel while the second one is the singularity due to the particle exchanged in the crossed process. The residues at $\beta = \frac{2\pi i}{3}$ are given by

$$\begin{aligned}
r_1 &= \text{Res}_{\beta=\frac{2\pi i}{3}} S_{00}^{11}(\beta) = 0 ; \\
r_2 &= \text{Res}_{\beta=\frac{2\pi i}{3}} S_{01}^{11}(\beta) = i \left(\frac{s\left(\frac{2}{5}\right)}{s\left(\frac{1}{5}\right)} \right)^2 \omega ; \\
r_3 &= \text{Res}_{\beta=\frac{2\pi i}{3}} S_{11}^{11}(\beta) = i \omega ;
\end{aligned} \tag{3.8}$$

$$\begin{aligned}
r_4 &= \text{Res}_{\beta=\frac{2\pi i}{3}} S_{11}^{01}(\beta) = i \left(\frac{s\left(\frac{2}{5}\right)}{s\left(\frac{1}{5}\right)} \right)^{\frac{1}{2}} \omega ; \\
r_5 &= \text{Res}_{\beta=\frac{2\pi i}{3}} S_{11}^{00}(\beta) = i \frac{s\left(\frac{2}{5}\right)}{s\left(\frac{1}{5}\right)} \omega ;
\end{aligned}$$

where

$$\omega = \frac{5}{9} \frac{s\left(\frac{1}{5}\right) s\left(\frac{1}{10}\right) s\left(\frac{4}{9}\right) s\left(\frac{1}{9}\right) s^2\left(\frac{5}{18}\right)}{s\left(\frac{3}{10}\right) s\left(\frac{1}{18}\right) s\left(\frac{7}{18}\right) s^2\left(\frac{2}{9}\right)} . \quad (3.9)$$

Their numerical values are collected in Table 1.

In the amplitude S_{00}^{11} there is no bound state in the direct channel but only the singularity coming from the state $|K_{11}\rangle$ exchanged in the t -channel. This is easily seen from Fig. 4 where we stretch the original amplitudes along the vertical direction (s -channel) and along the horizontal one (t -channel). Since the state $|K_{00}\rangle$ is not physical, the residue in the direct channel is zero. In the amplitude S_{01}^{11} we have the bound state $|K_{01}\rangle$ in the direct channel and the singularity due to $|K_{11}\rangle$ in the crossed channel. In S_{11}^{11} , the state $|K_{11}\rangle$ appears as a bound state in both channels. In S_{11}^{01} the situation is reversed with respect to that of S_{01}^{11} , as it should be from the crossing symmetry property (3.6): the state $|K_{11}\rangle$ appears in the t -channel and $|K_{01}\rangle$ in the direct channel. Finally, in S_{11}^{00} there is the bound state $|K_{11}\rangle$ in the direct channel but the residue on the t -channel pole is zero, again because $|K_{00}\rangle$ is unphysical. This situation is, of course, that obtained by applying crossing to S_{00}^{11} .

4 Energy levels, phase shifts and generalized statistics

The one-particle line a of fig. (1.a) corresponds to the state $|K_{11}\rangle$. This energy level is not doubly degenerate because the state $|K_{00}\rangle$ is forbidden by the RSOS selection rules, eq. (2.2). With periodic boundary conditions, the kink states $|K_{01}\rangle$ and $|K_{10}\rangle$ are projected out and $|K_{11}\rangle$ is the only one-particle state that can appear in the spectrum.

Consider the threshold line. On a strip with periodic boundary conditions we expect this energy level to be doubly degenerate. This because the states $|K_{01}K_{10}\rangle$ and

$|K_{10}K_{01}\rangle$ are identified under such boundary conditions whereas the state $|K_{11}K_{11}\rangle$ remains distinct. The conformal operators creating these states in the u.v. limit are $\Phi_{\frac{7}{16}, \frac{7}{16}}$ and $\Phi_{\frac{6}{10}, \frac{6}{10}}$, respectively (see lines marked b and c in fig. 1.a). We have checked that for large values of R these two lines approach each other faster than $1/r$ (as would be the case if the lowest line b was the only threshold line and c a line of momentum). However the truncation effects already present in this region prevents us from showing that they really approach each other exponentially³. In the following we assume that ideally these two levels are exponentially degenerate in the infrared limit.

The identification of the threshold lines described above holds only for the static configuration of two kinks with zero relative momentum. The situation is indeed different for the lines of momentum which approach the threshold. In fact, the S -matrix acting on $|K_{01}K_{10}\rangle$ can only produce $|K_{01}K_{10}\rangle$ as final state whereas $|K_{10}K_{01}\rangle$ and $|K_{11}K_{11}\rangle$ can mix through the processes of eq. (3.2). In order to determine the pattern of the energy levels obtained from TCSCA and to relate the scattering processes to the data of the original unperturbed CFT (along the line suggested in [12]), we would need a higher-level Bethe ansatz technique. This is because our actual situation deals with kink-like excitations in contrast to that of ref. [12] which considers only diagonal, breather-like S matrices. The Bethe-ansatz technique gets quite complicated in the case of a S -matrix with kink excitations. For the moment, it has been applied in few cases [13, 14]. In the light of these difficulties, we prefer here not to pursue such a program and instead concentrate on some properties of the phase shifts and generalized statistics which arises for kink excitations.

For real values of β , the amplitudes $S_{00}^{11}(\beta)$ and $S_{01}^{11}(\beta)$ are numbers of modulus 1. It

³Consider, for instance, figure 10 of ref. [2] (the case of low-temperature phase of thermal perturbation of the TIM). One observes that the onset of the exponential approach of degenerate excited levels usually occurs quite far from the value of R at which the two ground state energies coincide. As we discuss in sect 6, the situation of the subleading magnetic deformation of the TIM is even worse from a numerical point of view, because the anomalous dimension of the subleading magnetic operator is approximately $1/2$. In this case, it is therefore possible that the onset of the exponential approach of the higher levels takes place in a region of R strongly dominated by truncation effects.

is therefore convenient to define the following phase shifts

$$\begin{aligned} S_{00}^{11}(\beta) &\equiv e^{2i\delta_0(\beta)}; \\ S_{01}^{11}(\beta) &\equiv e^{2i\delta_1(\beta)}. \end{aligned} \quad (4.1)$$

The non-diagonal sector of the scattering processes is characterized by the 2×2 symmetric S -matrix

$$\begin{pmatrix} S_{11}^{11}(\beta) & S_{11}^{01}(\beta) \\ S_{11}^{01}(\beta) & S_{11}^{00}(\beta) \end{pmatrix}. \quad (4.2)$$

We can define the corresponding phase shifts by diagonalizing the matrix (4.2). The eigenvalues turn out to be the same functions in (4.1),

$$\begin{pmatrix} e^{2i\delta_0(\beta)} & 0 \\ 0 & e^{2i\delta_1(\beta)} \end{pmatrix}. \quad (4.3)$$

The phase shifts, for positive values of β , are shown in fig. 5. Asymptotically, they have the following limits

$$\begin{aligned} \lim_{\beta \rightarrow \pm\infty} e^{2i\delta_0(\beta)} &= e^{\pm \frac{6\pi i}{5}}; \\ \lim_{\beta \rightarrow \pm\infty} e^{2i\delta_1(\beta)} &= e^{\pm \frac{3\pi i}{5}}. \end{aligned} \quad (4.4)$$

There is a striking difference between the two phase shifts: while $\delta_1(\beta)$ is a monotonic decreasing function, starting from its value at zero energy $\delta_1(0) = \frac{\pi}{2}$, $\delta_0(\beta)$ shows a maximum for $\beta \sim \frac{\pi}{3}$ and then decreases to its asymptotic value $\frac{3\pi}{5}$. Its values are always larger than $\delta_0(0) = \frac{\pi}{2}$. Such different behaviour of the phase shifts is related to the presence of a zero very close to the real axis in the amplitude $e^{2i\delta_0(\beta)}$, i.e. at $\beta = i\frac{\pi}{9}$. This zero competes with the pole at $\beta = i\frac{\pi}{3}$ in creating a maximum in the phase shift. Similar behaviour also occurs in non-relativistic cases [10] and in the case of breather-like S matrices which contains zeros [11]. The presence of such a zero is deeply related to the absence of the pole in the s-channel of the amplitude $e^{2i\delta_0(\beta)}$. For the amplitude $e^{2i\delta_1(\beta)}$, the zero is located at $\beta = \frac{4\pi i}{9}$ (between the two poles) and therefore its contribution to the phase shift is damped with respect to that one coming from the poles. The net result is a monotonic decreasing phase shift.

Coming back to the 2×2 S -matrix of eq. (4.2), a basis of eigenvectors is given by

$$\begin{aligned} |\phi_1(\beta_1)\phi_1(\beta_2)\rangle &= A(\beta_{12}) (|K_{11}(\beta_1)K_{11}(\beta_2)\rangle + \chi_1(\beta_{12}) |K_{10}(\beta_1)K_{01}(\beta_2)\rangle) ; \\ |\phi_2(\beta_1)\phi_2(\beta_2)\rangle &= A(\beta_{12}) (|K_{11}(\beta_1)K_{11}(\beta_2)\rangle + \chi_2(\beta_{12}) |K_{10}(\beta_1)K_{01}(\beta_2)\rangle) . \end{aligned} \quad (4.5)$$

where $A(\beta_{12})$ is a normalization factor. In the asymptotic regime $\beta \rightarrow \infty$

$$\begin{aligned} \chi_1 &= - \frac{e^{-\frac{2\pi i}{5}} (a^2 + e^{\frac{6\pi i}{5}})}{a} \\ \chi_2 &= - \frac{e^{-\frac{2\pi i}{5}} (a^2 + e^{\frac{3\pi i}{5}})}{a} , \end{aligned} \quad (4.6)$$

and the probability P_{1001} to find a state $|K_{10}K_{01}\rangle$ in the vector $|\phi_2\phi_2\rangle$ w.r.t. the probability P_{1111} to find a state $|K_{11}K_{11}\rangle$ is given by the golden ratio

$$\frac{P_{1001}}{P_{1111}} = \frac{1}{a^2} = 2 \cos\left(\frac{\pi}{5}\right) . \quad (4.7)$$

For the state $|\phi_1\phi_1\rangle$, we have

$$\frac{P_{1001}}{P_{1111}} = a^2 = \frac{1}{2 \cos\left(\frac{\pi}{5}\right)} . \quad (4.8)$$

The “kinks” ϕ_1 and ϕ_2 have the generalized bilinear commutation relation [15, 16, 17]

$$\phi_i(t, x)\phi_j(t, y) = \phi_j(t, y)\phi_i(t, x) e^{2\pi i s_{ij} \epsilon(x-y)} . \quad (4.9)$$

The generalized “spin” s_{ij} is a parameter related to the asymptotic behaviour of the S -matrix. A consistent assignment is given by

$$\begin{aligned} s_{11} &= \frac{3}{5} = \frac{\delta_0(\infty)}{\pi} ; \\ s_{12} &= 0 ; \\ s_{22} &= \frac{3}{10} = \frac{\delta_1(\infty)}{\pi} . \end{aligned} \quad (4.10)$$

The implications of these generalized statistics will be discussed elsewhere. Here we only notice that, interesting enough, the previous monodromy properties are those of the chiral field $\Psi = \Phi_{\frac{6}{10},0}$ of the original CFT of the TIM. The operator product expansion of Ψ with itself reads

$$\Psi(z)\Psi(0) = \frac{1}{z^{\frac{6}{5}}} \mathbf{1} + \frac{C_{\Psi,\Psi,\Psi}}{z^{\frac{3}{5}}} \Psi(0) + \dots \quad (4.11)$$

where $C_{\Psi,\Psi,\Psi}$ is the structure constant of the OPE algebra. Moving z around the origin, $z \rightarrow e^{2\pi i}z$, the phase acquired from the first term on the right hand side of (4.11) comes from the conformal dimension of the operator Ψ itself. In contrast, the phase obtained from the second term is due to the insertion of an additional operator Ψ . A similar structure appears in the scattering processes of the “kinks” ϕ_i : in the amplitude of the kink ϕ_1 there is no bound state in the s-channel (corresponding to the “identity term” in (4.11)) whereas in the amplitude of ϕ_2 a kink can be created as a bound state for $\beta = \frac{2\pi i}{3}$ (corresponding to the “ Ψ term” in (4.11)). In the ultraviolet limit, the fields ϕ_i should give rise to the operator $\Psi(z)$, similarly to the case analyzed in [17]. The actual proof requires the analysis of the form factors and will be given elsewhere.

5 Zamolodchikov’s S -matrix for $\Phi_{2,1}$ perturbed TIM

The problem of finding a theoretical explanation for the energy levels of the $\Phi_{2,1}$ perturbed TIM was first discussed in a remarkable paper by Zamolodchikov [4]. In his notation, the one particle states are given by

$$|K_+\rangle, |K_-\rangle, |B\rangle, \quad (5.1)$$

which we can identify with our $|K_{10}\rangle$, $|K_{01}\rangle$ and $|K_{11}\rangle$, respectively. The two-particle amplitudes of the scattering processes were defined in [4] to be

$$\begin{aligned} |B(\beta_1)B(\beta_2)\rangle &= a(\beta_1 - \beta_2) |B(\beta_2)B(\beta_1)\rangle + b(\beta_1 - \beta_2) |K_+(\beta_2)K_-(\beta_1)\rangle \\ |K_-(\beta_1)B(\beta_2)\rangle &= c(\beta_1 - \beta_2) |K_-(\beta_2)B(\beta_1)\rangle \\ |B(\beta_1)K_+(\beta_2)\rangle &= c(\beta_1 - \beta_2) |B(\beta_2)K_+(\beta_1)\rangle \\ |K_-(\beta_1)K_+(\beta_2)\rangle &= d(\beta_1 - \beta_2) |K_-(\beta_2)K_+(\beta_1)\rangle \\ |K_+(\beta_1)K_-(\beta_2)\rangle &= e(\beta_1 - \beta_2) |K_+(\beta_2)K_-(\beta_1)\rangle + b(\beta_1 - \beta_2) |B(\beta_2)B(\beta_1)\rangle \end{aligned} \quad (5.2)$$

They are in correspondence with those of eq. (3.2) if we make the following assignments

$$a(\beta) \rightarrow S_{11}^{11}(\beta);$$

$$\begin{aligned}
b(\beta) &\rightarrow S_{11}^{10}(\beta) ; \\
c(\beta) &\rightarrow S_{01}^{11}(\beta) ; \\
d(\beta) &\rightarrow S_{00}^{11}(\beta) ; \\
e(\beta) &\rightarrow S_{11}^{00}(\beta) . .
\end{aligned} \tag{5.3}$$

In order to solve the Yang-Baxter equations which ensure the factorization of the scattering processes, Zamolodchikov noticed that the above amplitudes coincide with the definitions of the Boltzmann weights of the “Hard Square Lattice Gas”. Therefore, he borrowed Baxter’s solution [18] in the case where it reduces to trigonometric form

$$\begin{aligned}
a(\beta) &= \frac{\sin\left(\frac{2\pi}{5} + \lambda\beta\right)}{\sin\left(\frac{2\pi}{5}\right)} R(\beta) ; \\
b(\beta) &= e^{\delta\beta} \frac{\sin(\lambda\beta)}{\left[\sin\left(\frac{2\pi}{5}\right) \sin\left(\frac{\pi}{5}\right)\right]^{\frac{1}{2}}} R(\beta) ; \\
c(\beta) &= e^{-\delta\beta} \frac{\sin\left(\frac{\pi}{5} - \lambda\beta\right)}{\sin\left(\frac{\pi}{5}\right)} R(\beta) ; \\
d(\beta) &= e^{-2\delta\beta} \frac{\sin\left(\frac{\pi}{5} + \lambda\beta\right)}{\sin\left(\frac{\pi}{5}\right)} R(\beta) ; \\
e(\beta) &= e^{2\delta\beta} \frac{\sin\left(\frac{2\pi}{5} - \lambda\beta\right)}{\sin\left(\frac{2\pi}{5}\right)} R(\beta) .
\end{aligned} \tag{5.4}$$

Here δ and λ are arbitrary parameters and $R(\beta)$ is an arbitrary function. In order to fix completely the amplitudes, Zamolodchikov imposed the following requirements:

1. the unitarity conditions, eqs.(3.5);
2. the absence of a pole in the direct channel of the amplitude $d(\beta)$;
3. crossing symmetry, implemented in the following form

$$\begin{aligned}
a(\beta) &= a(i\pi - \beta) ; \\
b(\beta) &= c(i\pi - \beta) ; \\
d(\beta) &= e(i\pi - \beta) .
\end{aligned} \tag{5.5}$$

The final form of the S matrices is given by

$$\begin{aligned}
a(\beta) &= e^{-2i\pi\delta} \frac{\sin\left(\frac{2\pi-6i\beta}{5}\right) \sin\left(\frac{3\pi+6i\beta}{5}\right)}{\sin\left(\frac{\pi-6i\beta}{5}\right) \sin\left(\frac{2\pi+6i\beta}{5}\right)} ; \\
b(\beta) &= e^{-\delta(i\pi-\beta)} \frac{\sin\left(\frac{6i\beta}{5}\right) \sin\left(\frac{3\pi+6i\beta}{5}\right)}{\sin\left(\frac{\pi-6i\beta}{5}\right) \sin\left(\frac{2\pi+6i\beta}{5}\right)} ; \\
c(\beta) &= e^{-\delta\beta} \frac{\sin\left(\frac{\pi+6i\beta}{5}\right) \sin\left(\frac{3\pi+6i\beta}{5}\right)}{\sin\left(\frac{\pi-6i\beta}{5}\right) \sin\left(\frac{2\pi+6i\beta}{5}\right)} ; \\
d(\beta) &= e^{-2\delta\beta} \frac{\sin\left(\frac{3\pi+6i\beta}{5}\right)}{\sin\left(\frac{2\pi+6i\beta}{5}\right)} ; \\
e(\beta) &= e^{-2\delta(i\pi-\beta)} \frac{\sin\left(\frac{3\pi+6i\beta}{5}\right)}{\sin\left(\frac{\pi-6i\beta}{5}\right)} .
\end{aligned} \tag{5.6}$$

Herein δ is an imaginary number satisfying

$$e^{-2\pi i\delta} = \frac{s\left(\frac{1}{5}\right)}{s\left(\frac{2}{5}\right)} . \tag{5.7}$$

All amplitudes but $d(\beta)$ have a simple pole at $\beta = \frac{2\pi i}{3}$. Their residues are given by

$$\begin{aligned}
\tau_1 &= \text{Res}_{\beta=\frac{2\pi i}{3}} a(\beta) = i \frac{5}{6} \frac{\left(s\left(\frac{1}{5}\right)\right)^3}{\left(s\left(\frac{2}{5}\right)\right)^2} ; \\
\tau_2 &= \text{Res}_{\beta=\frac{2\pi i}{3}} b(\beta) = -i \frac{5}{6} \frac{\left(s\left(\frac{1}{5}\right)\right)^{\frac{13}{6}}}{\left(s\left(\frac{2}{5}\right)\right)^{\frac{7}{6}}} ; \\
\tau_3 &= \text{Res}_{\beta=\frac{2\pi i}{3}} c(\beta) = i \frac{5}{6} \frac{\left(s\left(\frac{1}{5}\right)\right)^{\frac{4}{3}}}{\left(s\left(\frac{2}{5}\right)\right)^{\frac{1}{3}}} ; \\
\tau_4 &= \text{Res}_{\beta=\frac{2\pi i}{3}} d(\beta) = 0 ; \\
\tau_5 &= \text{Res}_{\beta=\frac{2\pi i}{3}} e(\beta) = i \frac{5}{6} \frac{\left(s\left(\frac{1}{5}\right)\right)^{\frac{4}{3}}}{\left(s\left(\frac{2}{5}\right)\right)^{\frac{1}{3}}} .
\end{aligned} \tag{5.8}$$

Their numerical values are collected in Table 2.

In the asymptotic limit $\beta \rightarrow \infty$, all amplitudes but $a(\beta)$ have an oscillating behaviour

$$\begin{aligned}
a(\beta) &\sim e^{-2\pi i\delta} ; \\
b(\beta) &\sim e^{-\delta i\pi} e^{\delta\beta} e^{\frac{3\pi i}{5}} ;
\end{aligned}$$

$$\begin{aligned}
c(\beta) &\sim e^{-\delta\beta} e^{-\frac{3\pi i}{5}} ; \\
d(\beta) &\sim e^{-2\delta\beta} e^{-\frac{i\pi}{5}} ; \\
e(\beta) &\sim e^{-2\delta i\pi} e^{2\delta\beta} e^{\frac{-4\pi i}{5}} .
\end{aligned}
\tag{5.9}$$

Such oscillating behaviour was also found by the same author for a scattering model with Z_4 symmetry [19]. There, it was suggested that in the ultraviolet limit, the Z_4 theory has a limit cycle. Whether or not this is also the case for the S -matrix proposed by Zamolodchikov for the subleading magnetic perturbation of TIM, it is not clear to us that it is possible to match such oscillating behaviour to either a definite CFT in the ultraviolet limit, or a choice of generalized statistics for the kinks.

6 Finite-size effects

The study of the scaling region around a fixed point is simplified along those directions (in the space of coupling constants) which define an integrable massive field theory. In such cases, the knowledge of the theory on mass-shell (the S -matrix) makes it possible to characterize completely the dynamics even off-shell. In particular, using the Thermodynamical Bethe Ansatz method [20] one could compute the ground-state energy $E_0(R)$ for the theory on a cylinder of width R . If this computation could be performed for the subleading magnetic perturbation of TIM, it would become easy to decide which of the two proposed scattering theories is the correct one. Unfortunately, the Bethe-ansatz technique has not been extended to the case of an S -matrix with kink excitations. Only few examples have been worked out [13, 14]. However, we can get around this difficulty using the “Truncated Conformal Space Approach” (TCSA) [2, 3, 22] and the predictions of finite-size theory [4, 8].

The TCSA allows us to study the crossover from massless to massive behaviour in a theory with the space coordinate compactified on a circle of radius R . The method consists in truncating the infinite-dimensional Hilbert space of the CFT up to a level Λ in the Verma modules. Then, the off-critical Hamiltonian

$$H(\lambda, R) = H_0(R) + V(R) . \tag{6.1}$$

is numerically diagonalized. Here, $H_0(R)$ is the Hamiltonian of the fixed point [21]

$$H_0(R) = \frac{2\pi}{R} \left(L_0 + \overline{L}_0 - \frac{c}{12} \right) , \quad (6.2)$$

and $V(R)$ is the interaction term given by the perturbation

$$V(R) = \lambda \int_0^R \Phi_{r,s}(x) dx . \quad (6.3)$$

The matrix elements of $V(R)$ are computed in terms of the three-point functions of the scaling fields of the fixed point. An efficient algorithm has been developed for performing such computation [22]. In our case, the truncation is fixed at level 5 in the Verma modules. The parameter λ in (6.3) is a dimensionful coupling constant, related to the mass scale of the perturbed theory

$$[\lambda] = m^{2-2\Delta_{r,s}} . \quad (6.4)$$

In the following we fix the mass scale by $\lambda = 1$.

The energy levels $E_i(R)$ of the Hamiltonian (6.1) have the scaling form

$$E_i(R) = \frac{1}{R} F_i(mR) , \quad (6.5)$$

with the asymptotic behaviour

$$E_i(R) \simeq \begin{cases} \frac{2\pi}{R} \left(2\Delta_i - \frac{c}{12} \right) & , mR \ll 1 ; \\ \epsilon_0 m^2 R + M_i & , mR \gg 1 . \end{cases} \quad (6.6)$$

Here, Δ_i is the anomalous dimension of some scaling field in the ultraviolet regime and M_i is the (multi)particle mass term in the infrared limit. However, the above infrared asymptotic behaviour holds only in the ideal situation when the truncation parameter Λ goes to infinity. In practice, for finite Λ , the linear behaviour of eq. (6.6) is realized only within a finite region of the R axis. The large R behaviour is dictated by truncation effects. In order to find the physical regions, we make use of the following parameter (introduced in ref. [2])

$$\rho_i(R) = \frac{d \log E_i(R)}{d \log R} . \quad (6.7)$$

The parameter ρ_i is between the values $\rho_i = -1$ (in the ultraviolet region) and $\rho_i = 1$ (in the infrared one). In the limit of large R (the truncation-dominated regime), $\rho_i = 1 - 2\Delta$.

The “window” in R where the linear infrared behaviour holds depends upon the perturbing field and, for the case of operators with anomalous dimension $\Delta \geq \frac{1}{2}$, it can be completely shrunk away. This phenomenon is related to the divergences which appear in a perturbative expansion of the Hamiltonian (6.1), which must be renormalized. Under these circumstances, it is more convenient to consider the differences of energies, which are not renormalized.

In the case of the subleading magnetic perturbation of TIM, the anomalous dimension of $\Phi_{2,1}$ is $\Delta = \frac{7}{16}$, which is near $\frac{1}{2}$. Looking at fig. (1.a), we see that the onset of the infrared region of the two lowest levels is around $R \sim 2$ and persists only for few units in R . In this region one can check that they approach each other exponentially [2]

$$E_1 - E_0 \sim e^{-mR} , \quad (6.8)$$

and extract in this way the numerical mass of the kinks

$$m = 0.98 \pm 0.02 . \quad (6.9)$$

From fig. 6, we see that for the third level, that of one-particle state, the ultraviolet behaviour extends till $R \sim 0.5$. The crossover region is in the interval $0.5 \leq R \leq 2$. Beyond this interval, the infrared regime begins but the “window” of infrared behaviour is quite narrow and is in the vicinity of $R \sim 3$. In such small region, it becomes hard to extract any sensible result. To overcome this difficulty, it is better to consider the differences of energies with respect to those of the degenerate ground states. From fig. (1.b) one can easily read off the mass-gap and see that is consistent with the value extracted from the exponential approach of the two lowest levels. In fig. (1.b), the third line defines the threshold, with a mass-gap $2m$.

In the ideal situation $\Lambda \rightarrow \infty$, the crossover between the intermediate region ($mR \sim 1$) and the infrared one ($mR \gg 1$) is controlled by off-mass shell effects and has an exponential behaviour. The computation of these finite-size corrections has been put forward by Lüscher [8]. In the case of one-particle state, there are two leading off-mass-shell contributions coming from the processes shown in fig. 7. The first correction involves the on-mass-shell three-particle vertex Γ , which is extracted from the residue at

$\beta = \frac{2\pi i}{3}$ of the amplitudes $S_{11}^{11}(\beta)$ (in the case of RSOS S -matrix) and $a(\beta)$ (in the case of Zamolodchikov's S -matrix). The second correction comes from an integral over the momentum of the intermediate virtual particle, interacting via the S -matrix ($S_{11}^{11}(\beta)$ for the RSOS S -matrix and $a(\beta)$ for that of Zamolodchikov). Therefore we have

$$\begin{aligned} \Delta E(R) \equiv E_2(R) - E_0(R) = & m + i \frac{\sqrt{3}m}{2} \Gamma^2 \exp\left(-\frac{\sqrt{3}mR}{2}\right) \\ & - m \int_{-\infty}^{\infty} \frac{d\beta}{2\pi} e^{-mR \cosh \beta} \cosh \beta \left(S\left(\beta + \frac{i\pi}{2}\right) - 1 \right) . \end{aligned} \quad (6.10)$$

We have done the following. First we have computed numerically the integral on the intermediate particles in both cases of RSOS and Zamolodchikov's S -matrix and we have subtracted it from the numerical data obtained from the TCSA. After this subtraction, we have made a fit of the data with a function of the form

$$G(R) = A + B e^{-\frac{\sqrt{3}}{2}mR} + C e^{-mR} . \quad (6.11)$$

The first term should correspond to the mass term. The coefficient of second one is the quantity we need in order to extract the residue of the S -matrix at $\beta = \frac{2\pi i}{3}$

$$\frac{2}{\sqrt{3}m} B = i \operatorname{Res}_{\beta=\frac{2\pi i}{3}} \begin{cases} S_{11}^{11}(\beta) \\ a(\beta) \end{cases} . \quad (6.12)$$

The third term is a subleading correction related to the asymptotic approach of the lowest levels of our TCSA data to the theoretical vacuum energy $E_0(R)$.

In the case of RSOS S -matrix, the best fit gives the following values

$$\begin{aligned} A &= 0.97 \pm 0.02 ; \\ B &= -0.29 \pm 0.02 ; \\ C &= -0.36 \pm 0.02 . \end{aligned} \quad (6.13)$$

The corresponding curve is drawn in fig. 8, together with the data obtained from TCSA. The mass term agrees with our previous calculation (eq. (6.9)). The second term gives for the residue at $\beta = \frac{2\pi i}{3}$ the value 0.34 ± 0.02 . This is consistent with that of the RSOS S -matrix. In our fit procedure, the value of the residue we extracted through (6.12) is stable with respect to small variation of the mass value. Increasing (decreasing) m , B

increases (decreases) as well, in such a way that the residue takes the same value (into the numerical errors). This is a pleasant situation because it permits an iterative procedure for finding the best fit of the data: one can start with a trial value for m (let's say $m = 1$) and plug it into (6.11). From the A -term which comes out from the fit, one gets a new determination of the mass m that can be again inserted into (6.11) and so on. Continued iteration does not affect significantly the value we extract for the residue, but converges to an accurate measurement of the mass. The values in (6.13) were obtained in this way.

With Zamolodchikov's S -matrix the best fit of the data (with the same iterative procedure as before) gives the result

$$\begin{aligned} A &= 0.96 \pm 0.02 ; \\ B &= -1.10 \pm 0.02 ; \\ C &= 1.14 \pm 0.02 . \end{aligned} \tag{6.14}$$

The residue extracted from these data (1.29 ± 0.01) is not consistent with that one of the amplitude $a(\beta)$. The situation does not improve even if we *fix* the coefficient of $e^{-\frac{\sqrt{3}m}{2}}$ to be that one predicted by Table 2, namely $B = -0.158$ and leave as free parameters for a best fit A and C . In this case, our best determination of A and C were $A = 0.965$ and $C = -0.046$. The curve is plotted in fig. 8 together with the data obtained from TCSA.

7 Conclusions

The S -matrix proposed by Zamolodchikov for the subleading magnetic perturbation of TIM is a particular solution of the ‘‘Hard Square Lattice Gas’’ [18], fitted in such a way that it matches the physical picture coming from the TCSA data, i.e. the presence of two fundamental kinks and only one bound state thereof. But also the RSOS S -matrix computed in sect. 3 is a particular set of Boltzmann weights for the ‘‘Hard Square Lattice Gas’’ and it reproduces the same features. The key difference is how the crossing symmetry comes about: for the RSOS S -matrix, one finds a non-trivial charge conjugation matrix, eq. (3.6), whereas for Zamolodchikov's proposal the crossing symmetry occurs in a standard way, eq. (5.5). Their asymptotic behaviour for large real values of β is also

quite different: the RSOS S -matrix goes to a definite limit while the S -matrix proposed by Zamolodchikov is oscillating. For the RSOS case, the finite limit of the S -matrix allows us to introduce generalized statistics of the kink excitations [15, 16, 17].

The real discrimination between the two scattering theories proposed for the subleading magnetic deformation of TIM is seen by comparing them with a numerical “experiment”, i.e. from the study of the finite-size corrections of the energy levels obtained by the Truncated Conformal Space Approach. We have investigated this problem in sect. 6. The result suggests that the RSOS S -matrix gives a more appropriate description for the scattering processes of the massive excitations of the model. Though, the interesting question as to what kind of system the S -matrix of Zamolodchikov corresponds to, remains open.

Acknowledgments

We thank T. Miwa, A. Schwimmer and Al.B. Zamolodchikov for useful discussions. We are grateful to P. Orland for a critical reading of the paper. One of us (F.C.) thanks P. Di Vecchia for warm hospitality at NORDITA.

References

- [1] A.B. Zamolodchikov, *JETP Letters* **46** (1987),160; *Int. Journ. Mod. Phys.* **A3** (1988), 743; *Integrable field theory from CFT*, Proceeding of the Taniguchi Symposium, Kyoto 1988, to appear in *Advanced Studies in Pure Mathematics*.
- [2] M. Lässig, G. Mussardo and J.L. Cardy, *Nucl. Phys.* **B348** (1991), 591.
- [3] V.P. Yurov and A.B. Zamolodchikov, *Int. J. Mod. Phys.* **A5** (1990), 3221.
- [4] A.B. Zamolodchikov, *S-matrix of the Subleading Magnetic Perturbation of the Tricritical Ising Model*, PUTP 1195-90.
- [5] F.A. Smirnov, *Int. J. Mod. Phys.* **A6** (1991), 1407.
- [6] A.G. Izergin and V.E. Korepin, *Commun. Math. Phys.* **79** (1981), 303.
- [7] A.N. Kirillov and N. Yu. Reshetikhin, *Representations of the algebra $U_q(sl(2))$, q -orthogonal polynomials and invariants of links*, LOMI-preprint E-9-88.
- [8] M. Lüscher, *Comm. Math. Phys.* **104** (1986), 177; in *Progress in Gauge Field Theory* (Cargèse 1983), ed. G. 't Hooft et al., Plenum, New York 1984; in *Champs, Cordes, et Phénomènes Critiques*, proceedings of the 1988 Les Houches Summer School, ed. E. Brezin and J. Zinn-Justin, North Holland, Amsterdam, 1989.
- [9] G. Mussardo, *Integrable deformations of the non-unitary minimal conformal model $\mathcal{M}_{3,5}$* , NORDITA/91/54.
- [10] L.I. Schiff, *Quantum Mechanics*, McGraw-Hill, Inc., 1968.
- [11] F. Colomo and G. Mussardo, *unpublished*.
- [12] M. Lässig and M.J. Martins, *Nucl. Phys.* **B354** (1991), 666.
- [13] C.N. Yang, *Phys. Rev. Lett.* **19** (1967), 1312.
- [14] A.B. Zamolodchikov, *Nucl. Phys.* **B358** (1991), 497; *Nucl. Phys.* **B358** (1991), 524.

- [15] A. Swieca, *Fortschr. Phys.* **25** (1977), 303.
- [16] M. Karowski and H.J. Thun, *Nucl. Phys.* **B190** [FS3] (1981), 61.
- [17] F.A. Smirnov, *Comm. Math. Phys.* **132** (1990), 415.
- [18] R.J. Baxter, *Exactly Solved Models in Statistical Mechanics*, Academy Press, 1982.
- [19] A.B. Zamolodchikov, *Comm. Math. Phys.* **69** (1979), 165.
- [20] A.B. Zamolodchikov, *Nucl. Phys.* **B342** (1990), 695.
- [21] J.L. Cardy, *Nucl. Phys.* **270** [FS16] (1986), 186; *Nucl. Phys.* **B275** [FS17] (1986), 200; in *Champs, Cordes, et Phénomènes Critiques*, proceedings of the 1988 Les Houches Summer School, ed. E. Brezin and J. Zinn-Justin, North Holland, Amsterdam, 1989.
- [22] M. Lässig and G. Mussardo, *Computer Phys. Comm.* **66** (1991), 71.